

MULTIPLE SCLEROSIS SPECIALTY CARE PROGRAM

Phone: 888-216-1949 • Fax: 888-977-1391



©2017 KloudScript, Inc. - All rights reserved.

Name:	IATION:	2 PRESCRIBER INFORMATION:	
Name:		Name:	
Address:		Address:	
Oity:	State: Zip:	City: State: Z	
Phone:	Alt. Phone:	Phone: Fax:	
Email:		NPI: DEA:	
OOB: Gende	r: O M O F Caregiver:	Tax I.D.:	
Height: Weight:	Allergies:	Office Contact: Phone:	
3 STATEMENT OF I	MEDICAL NECESSITY: (Plea	se Attach All Medical Documentation)	
Date of Diagnosis:	ICD-10:	□ Relapse/Remitting	Progressive
		P ☐ Yes ☐ No ☐ Attach MRI Results Date:	
	raindication(s) to therapy? ☐ No ☐ Yes		
		rir res:	
II FIIOI AUGIOTIZAGOTI IS SCIII.	Bu. The Automatically Draft Appear of Flor	lew 4 Selful referred Formulary Atternatives	
4 INJECTION TRAIN	VING: O To Be Administered by Pharmacist	O Pharmacist to Provide Training O Patient Trained in MD Office O Manufa	acturer Nurse Support
5 PICK UP OR DEL	IVERY: O Delivery to Patient's F	Home O Delivery to Physician's Office O Pharmacy to	o Coordinate
6 INSURANCE INFO	PRMATION: Please Include From	nt and Back Copies of Pharmacy and Medical Card	
		choose both induction and maintenance dose where	annlicable)
Patient Name:	•	Patient's Date of Birth:	
Medication	Dosage & Strength	Direction	QTY Refills
□ AVONEX®	☐ 30mcg Prefilled Syringe☐ 30mcg Single Dose Vial☐ 30mcg Avonex Pen	☐ Inject 30mcg IM once a week☐ Titration: 7.5mcg weekly (over a 4 week period) until target dose is reached which is 30mcg	1 Kit
	30mcg Single Dose Vial	☐ Titration: 7.5mcg weekly (over a 4 week period) until target dose	1 Kit
□ AVONEX®	☐ 30mcg Single Dose Vial ☐ 30mcg Avonex Pen	☐ Titration: 7.5mcg weekly (over a 4 week period) until target dose is reached which is 30mcg ☐ Inject 0.25mg (1ml) SC every other day ☐ Titration: Weeks 1-2: Inject 0.0625mg/0.25ml SC every other day ☐ Weeks 3-4: Inject 0.125mg/0.50ml SC every other day ☐ Weeks 5-6: Inject 0.1875mg/0.75ml SC every other day	
□ AVONEX® □ BETASERON®	□ 30mcg Single Dose Vial □ 30mcg Avonex Pen □ 0.3mg Lyophilized Powder □ 20mg Prefilled Syringe	□ Titration: 7.5mcg weekly (over a 4 week period) until target dose is reached which is 30mcg □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: Inject 0.0625mg/0.25ml SC every other day Weeks 3-4: Inject 0.125mg/0.50ml SC every other day Weeks 5-6: Inject 0.1875mg/0.75ml SC every other day Weeks 7 and onward: Inject 0.25mg/1ml SC every other day □ Inject 20mg SC daily □ Inject 40mg SC three times per week □ Other □ Inject 0.25mg (1ml) SC every other day	1 Kit
□ AVONEX® □ BETASERON®	□ 30mcg Single Dose Vial □ 30mcg Avonex Pen □ 0.3mg Lyophilized Powder □ 20mg Prefilled Syringe	□ Titration: 7.5mcg weekly (over a 4 week period) until target dose is reached which is 30mcg □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: Inject 0.0625mg/0.25ml SC every other day Weeks 3-4: Inject 0.125mg/0.50ml SC every other day Weeks 5-6: Inject 0.1875mg/0.75ml SC every other day Weeks 7 and onward: Inject 0.25mg/1ml SC every other day □ Inject 20mg SC daily □ Inject 40mg SC three times per week □ Other	1 Kit
□ AVONEX® □ BETASERON® □ COPAXONE®	□ 30mcg Single Dose Vial □ 30mcg Avonex Pen □ 0.3mg Lyophilized Powder □ 20mg Prefilled Syringe □ 40mg Prefilled Syringe	□ Titration: 7.5mcg weekly (over a 4 week period) until target dose is reached which is 30mcg □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: Inject 0.0625mg/0.25ml SC every other day Weeks 3-4: Inject 0.125mg/0.50ml SC every other day Weeks 5-6: Inject 0.1875mg/0.75ml SC every other day Weeks 7 and onward: Inject 0.25mg/1ml SC every other day □ Inject 20mg SC daily □ Inject 40mg SC three times per week □ Other □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: 0.0625mg/0.25ml SC every other day Weeks 3-4: 0.125mg/0.50ml SC every other day Weeks 5-6: 0.1875mg/0.75ml SC every other day	1 Kit
□ AVONEX® □ BETASERON® □ COPAXONE® □ EXTAVIA®	□ 30mcg Single Dose Vial □ 30mcg Avonex Pen □ 0.3mg Lyophilized Powder □ 20mg Prefilled Syringe □ 40mg Prefilled Syringe □ 0.3mg Lyophilized Powder	□ Titration: 7.5mcg weekly (over a 4 week period) until target dose is reached which is 30mcg □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: Inject 0.0625mg/0.25ml SC every other day Weeks 3-4: Inject 0.125mg/0.50ml SC every other day Weeks 5-6: Inject 0.1875mg/0.75ml SC every other day Weeks 7 and onward: Inject 0.25mg/1ml SC every other day □ Inject 20mg SC daily □ Inject 40mg SC three times per week □ Other □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: 0.0625mg/0.25ml SC every other day Weeks 3-4: 0.125mg/0.50ml SC every other day Weeks 5-6: 0.1875mg/0.75ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day □ Take one capsule by mouth once daily	1 Kit
□ AVONEX® □ BETASERON® □ COPAXONE® □ EXTAVIA® □ GILENYA®	□ 30mcg Single Dose Vial □ 30mcg Avonex Pen □ 0.3mg Lyophilized Powder □ 20mg Prefilled Syringe □ 40mg Prefilled Syringe □ 0.3mg Lyophilized Powder □ 0.5mg Capsule	□ Titration: 7.5mcg weekly (over a 4 week period) until target dose is reached which is 30mcg □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: Inject 0.0625mg/0.25ml SC every other day Weeks 3-4: Inject 0.125mg/0.50ml SC every other day Weeks 5-6: Inject 0.1875mg/0.75ml SC every other day Weeks 7 and onward: Inject 0.25mg/1ml SC every other day □ Inject 20mg SC daily □ Inject 40mg SC three times per week □ Other □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: 0.0625mg/0.25ml SC every other day Weeks 3-4: 0.125mg/0.50ml SC every other day Weeks 5-6: 0.1875mg/0.75ml SC every other day □ Take one capsule by mouth once daily □ Other	1 Kit 1 Kit
□ AVONEX® □ BETASERON® □ COPAXONE® □ EXTAVIA® □ GILENYA®	□ 30mcğ Single Dose Vial □ 30mcg Avonex Pen □ 0.3mg Lyophilized Powder □ 20mg Prefilled Syringe □ 40mg Prefilled Syringe □ 0.3mg Lyophilized Powder □ 0.5mg Capsule □ 20mg Prefilled Syringe	□ Titration: 7.5mcg weekly (over a 4 week period) until target dose is reached which is 30mcg □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: Inject 0.0625mg/0.25ml SC every other day Weeks 3-4: Inject 0.125mg/0.50ml SC every other day Weeks 5-6: Inject 0.1875mg/0.75ml SC every other day Weeks 7 and onward: Inject 0.25mg/1ml SC every other day □ Inject 20mg SC daily □ Inject 40mg SC three times per week □ Other □ Inject 0.25mg (1ml) SC every other day Weeks 3-4: 0.125mg/0.25ml SC every other day Weeks 3-4: 0.125mg/0.50ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day □ Take one capsule by mouth once daily □ Other □ Inject 20mg SC daily	1 Kit 1 Kit
□ AVONEX® □ BETASERON® □ COPAXONE® □ EXTAVIA® □ GILENYA® □ GLATOPA™	□ 30mcğ Single Dose Vial □ 30mcg Avonex Pen □ 0.3mg Lyophilized Powder □ 20mg Prefilled Syringe □ 40mg Prefilled Syringe □ 0.3mg Lyophilized Powder □ 0.5mg Capsule □ 20mg Prefilled Syringe □ Titration Pack □ 22mcg Prefilled Syringe □ 44mcg Prefilled Syringe □ Rebidose® 22mcg Autoinjector	□ Titration: 7.5mcg weekly (over a 4 week period) until target dose is reached which is 30mcg □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: Inject 0.0625mg/0.25ml SC every other day Weeks 3-4: Inject 0.125mg/0.50ml SC every other day Weeks 5-6: Inject 0.1875mg/0.75ml SC every other day Weeks 7 and onward: Inject 0.25mg/1ml SC every other day Weeks 7 and onward: Inject 0.25mg/1ml SC every other day Inject 40mg SC three times per week □ Other □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: 0.0625mg/0.25ml SC every other day Weeks 3-4: 0.125mg/0.50ml SC every other day Weeks 5-6: 0.1875mg/0.75ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day □ Take one capsule by mouth once daily □ Other □ Inject 20mg SC daily □ Titration Pack Rebidose (six 8.8mcg pre-filled autoinjectors and six 22 mcgpre-filled autoinjectors) □ For 22mcg SC 3 times per week maintenance dose: ■ Weeks 1 & 2: Inject 4.4mcg 3 times per week ■ Weeks 5 and onward: Inject 22mcg 3 times per week ■ Weeks 5 and onward: Inject 22mcg 3 times per week ■ Weeks 1 & 2: Inject 8.8mcg 3 times per week ■ Weeks 3 & 4: Inject 8.8mcg 3 times per week ■ Weeks 3 & 4: Inject 8.8mcg 3 times per week	1 Kit 1 Kit 1 Kit
□ AVONEX® □ BETASERON® □ COPAXONE® □ EXTAVIA® □ GILENYA® □ GLATOPA™ □ REBIF®	□ 30mcğ Single Dose Vial □ 30mcg Avonex Pen □ 0.3mg Lyophilized Powder □ 20mg Prefilled Syringe □ 40mg Prefilled Syringe □ 0.3mg Lyophilized Powder □ 0.5mg Capsule □ 20mg Prefilled Syringe □ Titration Pack □ 22mcg Prefilled Syringe □ 44mcg Prefilled Syringe □ 44mcg Prefilled Syringe □ Rebidose® 22mcg Autoinjector □ Rebidose® 44mcg Autoinjector	□ Titration: 7.5mcg weekly (over a 4 week period) until target dose is reached which is 30mcg □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: Inject 0.0625mg/0.25ml SC every other day Weeks 3-4: Inject 0.125mg/0.50ml SC every other day Weeks 5-6: Inject 0.1875mg/0.75ml SC every other day Weeks 7 and onward: Inject 0.25mg/1ml SC every other day Weeks 7 and onward: Inject 0.25mg/1ml SC every other day Inject 40mg SC three times per week □ Other □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: 0.0625mg/0.25ml SC every other day Weeks 3-4: 0.125mg/0.50ml SC every other day Weeks 5-6: 0.1875mg/0.75ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day □ Take one capsule by mouth once daily □ Other □ Inject 20mg SC daily □ Titration Pack Rebidose (six 8.8mcg pre-filled autoinjectors and six 22 mcgpre-filled autoinjectors) □ For 22mcg SC 3 times per week maintenance dose: ■ Weeks 1 & 2: Inject 4.4mcg 3 times per week ■ Weeks 5 and onward: Inject 22mcg 3 times per week ■ Weeks 5 and onward: Inject 22mcg 3 times per week ■ Weeks 1 & 2: Inject 8.8mcg 3 times per week ■ Weeks 3 & 4: Inject 8.8mcg 3 times per week ■ Weeks 3 & 4: Inject 8.8mcg 3 times per week	1 Kit 1 Kit 1 Kit 30